

NO-A187 742

THEORY OF TWO-PHOTON EMISSION FROM ATOMIC INNER SHELLS
(U) OREGON UNIV EUGENE D GUO JUL 87 AFOSR-TR-87-1424
AFOSR-87-8826

1/1

UNCLASSIFIED

F/G 28/5

NL



REPORT DOCUMENTATION PAGE

AD-A187 742 JLE		1b RESTRICTIVE MARKINGS	
		3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release DISRTIBUTION UNLIMITED	
4 PERFORMING ORGANIZATION REPORT NUMBER(S)		5 MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 87-1424	
6a NAME OF PERFORMING ORGANIZATION UNIVERSITY OF OREGON DEPARTMENT OF PHYSICS	6b OFFICE SYMBOL (if applicable)	7a NAME OF MONITORING ORGANIZATION AFOSR/NP	
6c ADDRESS (City, State, and ZIP Code) EUGENE, OR 97403		7b ADDRESS (City, State, and ZIP Code) Bldg 410 BOLLING AFB, DC 20332	
8a NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR	8b OFFICE SYMBOL (if applicable) NP	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER CRANT #AFOSR-87-0026	
8c ADDRESS (City, State, and ZIP Code) Bldg 410 BOLLING AFB DC 20332-6418		10 SOURCE OF FUNDING NUMBERS PROGRAM ELEMENT NO. 1611001 PROJECT NO. 2301 TASK NO. A4 WORK UNIT NO. ACCESSION NO.	
11 TITLE (Include Security Classification) THEORY OF TWO-PHOTON EMISSION FROM ATOMIC INNER SHELLS			
12 PERSONAL AUTHOR(S) DONG-SHENG GUO			
13a TYPE OF REPORT PREPRINT	13b TIME COVERED FROM 15 JAN 87 TO 15 JAN 88	14 DATE OF REPORT (Year, Month, Day) JULY 1987	15 PAGE COUNT 16
16 SUPPLEMENTARY NOTATION SUBMITTED TO PHYSICAL REVIEW A			
17 COSATI CODES FIELD GROUP SUB-GROUP		18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number) MULTIPHOTON PROCESSES	
19 ABSTRACT (Continue on reverse if necessary and identify by block number) See attached			
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21 ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a NAME OF RESPONSIBLE INDIVIDUAL DR. BARRY Z. KELLEY		22b TELEPHONE (Include Area Code) (202) 767-4408	
		22c OFFICE SYMBOL NP	

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted
All other editions are obsolete

SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

87 10 14 389

DTIC
ELECTE
NOV 16 1987
S H D

~~To be~~ submitted to
Physical Review A

Theory of two-photon emission from atomic inner shells

Dong-Sheng Guo

Department of Physics and Chemical Physics Institute

University of Oregon

Eugene, Oregon 97403

We prove from quantum electrodynamics that two-photon emission in transitions between atomic inner shells can be treated in lowest-order approximation in the single-particle model. By taking many-body effects into account ^{to} we show that, in the reduced single-particle model, the Pauli exclusion principle does not prohibit summing over all possible intermediate states, including core states. Some transition-rate formulas are presented that are useful for numerical calculations.

I. INTRODUCTION

Recent developments in experimental techniques have made it possible to measure two-photon transitions between atomic inner-shell vacancy states.¹ The results must be interpreted relativistically. Existing theoretical work has been based on the hydrogenic²⁻⁴ or nonrelativistic self-consistent-field (SCF)⁴⁻⁶ models, except for the recent relativistic SCF calculation of Mu and Crasemann.⁷ The latter work suggests exploration of some interesting questions: Is the Pauli exclusion principle violated if one includes occupied orbitals among the intermediate states, as done in Ref. 7? What are the theoretical similarities and differences between two-photon processes in hydrogenic vs. inner-shell atomic systems? In this paper we address these questions. In Sec. II, we analyze the theory of two-photon transitions on the basis of quantum electrodynamics (QED). The possible intermediate states are identified automatically; they can be classified into two types according to time ordering. One type is advanced: the electron final state occurs when the intermediate states do, which is in advance of the disappearance of the initial state. The second type is retarded: the time order for the electron is first initial state, then intermediate states, then final state. In this derivation, the Pauli exclusion principle is built in by commutation relations of operators. The final expres-



For	
I	<input checked="" type="checkbox"/>
d	<input type="checkbox"/>
ion	<input type="checkbox"/>

ity Codes
and/or

Special

A-1

sion for the S-matrix element shows that the problem can be formally treated as a time-independent one, with the electron moving in the field of the nucleus and the other electrons. In the reduced model, since there is no time-order difference, the original two types of intermediate states patch up a new set of intermediate states which is a complete set. The Pauli principle is thus seen to allow summation over all possible intermediate states, including occupied inner-shell states.

In Sec. III, a time-independent many-body approach is delineated. In Sec. IV, as an application of the results of earlier sections, we provide formulas for matrix elements, transition rates, and differential equations to be solved, which are formally the same as for the hydrogenic case. The only difference is the external potential acting on the electron. It is not that an inner-shell vacancy is trivially equivalent to the hydrogenic case: there are many-body effects in inner shells, particularly the restrictions imposed by the Pauli exclusion principle, while the hydrogenic system consists of just a single bound electron. Nor should the reader think that this paper's result is just a simple repetition of the conclusions of Brown et al.⁸ regarding gamma-ray scattering, which are based on a single-electron, i.e. hydrogenic, model. If we consider many-body effects and the built-in Pauli exclusion principle in inner-shell gamma-ray elastic scattering, the core contributions should in fact be excluded when summing over intermediate states (unlike the procedure in Ref. 8). By contrast, as shown here, all intermediate states must be included (as done in Ref. 7) when one

calculates two-photon inner-shell transition probabilities, taking the Pauli exclusion principle into account.

II. QUANTUM-ELECTRODYNAMIC APPROACH AND THE QUESTION OF INTERMEDIATE STATES

The dominant part of two-photon emission is of the double electric-dipole (2E1) type; this is the only kind considered here.

In the theory of QED, the second-order scattering operator S can be expressed in Furry's representation⁹ as

$$S = -\frac{\alpha}{2} \iint d^4x d^4x' T(j^\mu(x) j^\nu(x') A_\mu(x) A_\nu(x')) \quad (1)$$

The matrix element is

$$S_{21} = \langle 0 | \langle \text{core} | c_1 c_2 a_1^\dagger S a_1 | \text{core} \rangle | 0 \rangle \quad (2)$$

where $|\text{core}\rangle$, in the lowest-order approximation, stands for an antisymmetrically filled core wave function, and $|0\rangle$, for a photon vacuum state. We denote photon creation and annihilation operators by c^\dagger and c , respectively, and electron creation and annihilation operators by a^\dagger and a . The remainder of the notation is of standard usage. Natural units ($\hbar=c=1$) are employed throughout. $\alpha = 1/137.036$.

The electron field operator can be expressed as

$$\begin{aligned} \psi(t, \underline{r}) &= \sum_{\underline{a}} a_{\underline{a}} \psi_{\underline{a}}(\underline{r}) \exp(-i\varepsilon_{\underline{a}} t) + \sum_{\underline{n}} a_{\underline{n}} \psi_{\underline{n}}(\underline{r}) \exp(-i\varepsilon_{\underline{n}} t) \\ \bar{\psi}(t, \underline{r}) &= \sum_{\underline{a}} a_{\underline{a}}^\dagger \bar{\psi}_{\underline{a}}(\underline{r}) \exp(i\varepsilon_{\underline{a}} t) + \sum_{\underline{n}} a_{\underline{n}}^\dagger \bar{\psi}_{\underline{n}}(\underline{r}) \exp(i\varepsilon_{\underline{n}} t) \end{aligned} \quad (3)$$

The summation index a runs over all core states, and n , over all the excited states.

After time-ordered contraction manipulation, the S matrix element takes the form

$$S_{21} = -i\alpha \iint d^4x d^4x' \bar{\psi}_1(x') \gamma^\nu G(x, x') \gamma^\mu \psi_2(x) (A_{1\mu}^* A_{2\nu}' + A_{2\mu}^* A_{1\nu}') \quad (4)$$

where

$$G(x, x') = -i \langle \text{core} | T \bar{\psi}(x) \psi(x') | \text{core} \rangle \quad (5)$$

is the propagator for an electron hole, and $\psi_1(x)$ and $\psi_2(x)$ are bispinor wave functions. The indices 1 and 2 denote the initial and final states of electrons, respectively.

The explicit form for the electron hole propagator can be shown to be

$$G(x, x') = \frac{1}{2\pi} \left[\sum_{a \in f} \int_{-\infty}^{+\infty} d\varepsilon \frac{\psi_a(x') \bar{\psi}_a(x)}{\varepsilon + \varepsilon_a + i\eta} e^{-i\varepsilon(t-t')} - \sum_{n \in f} \int_{-\infty}^{+\infty} d\varepsilon \frac{\psi_n(x') \bar{\psi}_n(x)}{\varepsilon - \varepsilon_n + i\eta} e^{-i\varepsilon(t'-t)} \right] \quad (6)$$

The first term in the bracket on the right-hand side of Eq. (6), which involves core intermediate states, is an advanced Green's function which describes the $t > t'$ process; the second term which involves excited intermediate states, is a retarded Green's function which describes the $t' > t$ process. This

distinction is important in understanding the physical meaning of the two types of intermediate states.

The two-photon processes can be described by Feynman diagrams. We specify the initial and final hole states in the following examples. Time progresses from left to right. A solid line with arrow opposite to the time direction represents an electron hole, a line with arrow in the time direction represents a real electron. There are two equivalent ways of expressing the advanced process, as shown in Fig. 1. Diagram 1(a) describes the process in terms of a hole, and diagram 1(b), in terms of an electron. For the retarded process, there are two similar diagrams, shown in Fig. 2.

From this analysis, we obtain only two types of intermediate states. Two other types of intermediate states could in principle be constructed, but do not occur in our derivation. One of these pertains to retarded real core states; the disappearance of this type embodies the Pauli exclusion principle. The other type of intermediates that does not appear here involves advanced excited hole states; this type is excluded by the fact that an excited state cannot be annihilated from the core.

Electromagnetic waves and electron waves have the forms

$$A_1(x) = A_1(\underline{r}) e^{-i\omega_1 t}, \quad A_2(x) = A_2(\underline{r}) e^{-i\omega_2 t};$$

$$\psi_1(x) = \psi_1(\underline{r}) e^{-i\varepsilon_1 t}, \quad \psi_2(x) = \psi_2(\underline{r}) e^{-i\varepsilon_2 t}.$$
(7)

respectively. After we integrate over time t , then t' , and then

energy , we find the following expression for the S matrix element:

$$S_{21} = -i 2\pi\alpha \int d^3\underline{r} \int d^3\underline{r}' \bar{\Psi}_1(\underline{r}') \gamma^\nu$$

$$\times \left[\sum_a \frac{\Psi_a(\underline{r}') \bar{\Psi}_a(\underline{r})}{\omega_1 - \varepsilon_2 + \varepsilon_a + i\eta} A_{1\mu}^*(\underline{r}) A_{2\nu}^*(\underline{r}') - \sum_n \frac{\Psi_n(\underline{r}') \bar{\Psi}_n(\underline{r})}{\varepsilon_1 + \omega_2 - \varepsilon_n + i\eta} A_{1\mu}^*(\underline{r}) A_{2\nu}^*(\underline{r}') \right. \quad (8)$$

$$\left. + \sum_a \frac{\Psi_a(\underline{r}') \bar{\Psi}_a(\underline{r})}{\omega_2 - \varepsilon_2 + \varepsilon_a + i\eta} A_{2\mu}^*(\underline{r}) A_{1\nu}^*(\underline{r}') - \sum_n \frac{\Psi_n(\underline{r}') \bar{\Psi}_n(\underline{r})}{\varepsilon_1 + \omega_1 - \varepsilon_n + i\eta} A_{2\mu}^*(\underline{r}) A_{1\nu}^*(\underline{r}') \right] \gamma^\mu \Psi_2(\underline{r}) \delta(\varepsilon_2 - \varepsilon_1 - \omega_1 - \omega_2)$$

The δ -function factor expresses the energy conservation law

$$\varepsilon_2 - \varepsilon_1 = \omega_1 + \omega_2 \quad (9)$$

By using this energy conservation condition, we can rewrite the energy denominators. It then becomes possible to combine the two types of Green's function into one with a complete set of wave functions as its intermediate states:

$$S_{21} = -i 2\pi\alpha \iint d^3\underline{r} d^3\underline{r}' \bar{\Psi}_1(\underline{r}') \gamma^\nu$$

$$\times \left[\sum_i \frac{\Psi_i(\underline{r}') \bar{\Psi}_i(\underline{r})}{\varepsilon_i + \omega_1 - \varepsilon_2} A_{1\mu}^*(\underline{r}) A_{2\nu}^*(\underline{r}') + \sum_i \frac{\Psi_i(\underline{r}') \bar{\Psi}_i(\underline{r})}{\varepsilon_i + \omega_2 - \varepsilon_2} A_{2\mu}^*(\underline{r}) A_{1\nu}^*(\underline{r}') \right] \quad (10)$$

$$\times \gamma^\mu \Psi_2(\underline{r}) \delta(\varepsilon_2 - \varepsilon_1 - \omega_1 - \omega_2)$$

Here, the index i runs over all possible intermediate states.

The expression for the S matrix element is thus reduced to a time-independent single-particle version, which is formally the same as in the hydrogenic model if we call the electron 2 as the initial state, the electron 1 as the final state.

The only differences between the two models are in two aspects: (1) the hydrogenic electron moves in the Coulomb field of the nucleus, while the electron displaced by the inner-shell hole moves in the combined field of the nucleus and the remaining electrons. (2) The summing and averaging over initial and final states, respectively, are opposite in the inner-shell case to that of outer shells. We have thus proved a theorem which is useful in deriving formulas for numerical calculations:

THEOREM I.

Two-E1 photon emission from atomic inner shells, filling one vacancy, can be expressed in the lowest-order approximation in a single-particle formulation which is formally the same as in the hydrogenic case, with the electron hole being treated as a real electron. The only differences are in the potential acting on the electron and the treatment of summing and averaging over the initial and final states.

III. TIME-INDEPENDENT MANY-BODY APPROACH

In a time-independent many-body approach, the initial and final states can be written as

$$|1\rangle = a_1 |core\rangle, \quad |2\rangle = a_2 |core\rangle \quad (11)$$

As discussed in Sec. II, we have two types of intermediate states, advanced and retarded. In time-independent many-body perturbation theory, these can be written

$$a_2 |core\rangle \quad (12)$$

and

$$a_n^\dagger a_s a_i | \text{core} \rangle \quad (13)$$

The dipole operator has the form

$$Q^m = \sum_{ij} q_{ij}^m a_i^\dagger a_j \quad (14)$$

where m is the magnetic quantum number. The time-independent many-body expression for the emission rate is

$$\frac{dW}{d\omega_1} = \omega_1^3 \omega_2^3 \frac{8\alpha^2}{9\pi} \sum_{m,m_2} \left| \sum_i \frac{\langle 2 | Q^{m_2} | i \rangle \langle i | Q^{m_1} | 1 \rangle}{E_i - E_i - \omega_1} + \sum_i \frac{\langle 2 | Q^{m_1} | i \rangle \langle i | Q^{m_2} | 1 \rangle}{E_i - E_i - \omega_2} \right|^2 \quad (15)$$

After contractions, this form is reduced to a single-body form

$$\frac{dW}{d\omega_1} = \omega_1^3 \omega_2^3 \frac{8\alpha^2}{9\pi} \sum_{m,m_2} \left| \sum_a \frac{q_{a2}^{m_2} q_{1a}^{m_1}}{E_a - E_1 + E_2 - \omega_2} + \sum_a \frac{q_{a2}^{m_1} q_{1a}^{m_2}}{E_a - E_1 + E_2 - \omega_2} - \sum_n \frac{q_{1n}^{m_2} q_{n2}^{m_1}}{E_2 - E_n - \omega_1} - \sum_n \frac{q_{1n}^{m_1} q_{n2}^{m_2}}{E_2 - E_n - \omega_1} \right|^2 \quad (16)$$

The many-body forms of the two types of intermediate states are now reduced to core states and excitations in the single-body form. If we impose the energy conservation law Eq. (9), the Green's function that involves core states and the one that involves excitations are combined into a new Green's function that involves a complete set of intermediate states. This proves Theorem I over again. We therefore have the final form for the emission rate:

$$\frac{dW}{d\omega_1} = \omega_1^3 \omega_2^3 \frac{8\alpha^2}{9\pi} \sum_{m,m_2} \left| \sum_i \frac{q_{1i}^{m_2} q_{i2}^{m_1}}{E_i - E_2 + \omega_1} + \sum_i \frac{q_{1i}^{m_1} q_{i2}^{m_2}}{E_i - E_2 + \omega_2} \right|^2 \quad (17)$$

If we specify the initial state for the atom as a $[1s]$ hole state, the possible final states will be $[2s]$, $[3s]$, $[3d_{3/2}]$, $[3d_{5/2}]$, $[4s]$, . . . , hole states. The possible intermediate states are of two types. The ones derived from retardation are of the type of single-hole p states of the core, and the ones derived from advancement are of the type of the states with one excited p electron and two holes in the core. This agrees with the classification of Aberg,⁵ and of Freund and Barnett.⁶ E.J. Robinson,¹⁰ M. Crance and M. Aymar¹¹ had similar treatments and conclusions for core excitations in outer shell problems.

The time-independent many-body approach leads to the same result as the QED approach. It is simpler, but involves some imposed conditions, such as those regarding the intermediate states and the energy conservation law. It cannot reveal the time-ordered process, and hence leads to less physical insight.

IV. MATRIX ELEMENTS AND EMISSION RATES

Theorem I reduces the inner-shell vacancy two-photon decay problem to one tractable in the single-particle model. The formulas for the hydrogenic case can all be adapted to the many-body problem, but the electron wave functions must be calculated for different potentials. A formalism for relativistic two-photon emission in the hydrogenic case has been given by Goldman

and Drake.⁷ We follow theirs and Tabak et al.⁸ which is equivalent to the approach of Kruer et al.⁸ After integrating over the angular distribution of the scattered electron, summing over all possible final states and summing over all possible initial states, the 2E1 decay rates per unit time in differential form are as follows:

$s \rightarrow s$

$$\begin{aligned} \frac{dW}{d\omega_1} = \frac{8\alpha^2}{27\pi} \omega_1^3 \omega_2^3 \left\{ \frac{1}{3} [E_1^2(\omega_1, \omega_2) + E_1^2(\omega_2, \omega_1)] + \frac{2}{3} [E_2^2(\omega_1, \omega_2) + E_2^2(\omega_2, \omega_1)] \right. \\ \left. - \frac{2}{9} E_1(\omega_1, \omega_2) E_1(\omega_2, \omega_1) + \frac{4}{9} E_{-2}(\omega_1, \omega_2) E_{-2}(\omega_2, \omega_1) \right. \\ \left. + \frac{8}{9} [E_{-2}(\omega_1, \omega_2) E_1(\omega_2, \omega_1) + E_1(\omega_1, \omega_2) E_{-2}(\omega_2, \omega_1)] \right\} \end{aligned}$$

$d_{3/2} \rightarrow s$

$$\begin{aligned} \frac{dW}{d\omega_1} = \frac{8\alpha^2}{27\pi} \omega_1^3 \omega_2^3 \left\{ \frac{2}{3} [E_1^2(\omega_1, \omega_2) + E_1^2(\omega_2, \omega_1)] + \frac{2}{15} [E_{-2}^2(\omega_1, \omega_2) + E_{-2}^2(\omega_2, \omega_1)] \right. \\ \left. + \frac{8}{9} E_1(\omega_1, \omega_2) E_1(\omega_2, \omega_1) - \frac{8}{45} E_{-2}(\omega_1, \omega_2) E_{-2}(\omega_2, \omega_1) \right. \\ \left. + \frac{4}{9} [E_{-2}(\omega_1, \omega_2) E_1(\omega_2, \omega_1) + E_1(\omega_1, \omega_2) E_{-2}(\omega_2, \omega_1)] \right\} \end{aligned}$$

$$d_{5/2} \rightarrow s$$

$$\frac{dW}{d\omega_1} = \frac{8\alpha^2}{27\pi} \omega_1^3 \omega_2^3 \left\{ \frac{6}{5} [E_{-2}^2(\omega_1, \omega_2) + E_{-2}^2(\omega_2, \omega_1)] + \frac{12}{5} E_{-2}(\omega_1, \omega_2) E_{-2}(\omega_2, \omega_1) \right\} \quad (20)$$

The transition amplitude is given by

$$E_K(\omega_1, \omega_2) = \frac{3}{\omega_2} \int_0^\infty dr \left[S_{KK'}(r, \omega_1) U_K(r, \omega_2) + T_{KK'}(r, \omega_1) V_K(r, \omega_2) \right] \quad (21)$$

where $S_{KK'}$ and $T_{KK'}$ are solutions of the perturbed Dirac equation

$$\begin{aligned} (m - \varepsilon_1 + \omega_1 + V(r)) S_{KK'}(r, \omega_1) + \left(\frac{d}{dr} - \frac{\kappa}{r} \right) T_{KK'}(r, \omega_1) &= \frac{3}{\omega_1} K_{KK'}(r, \omega_1) \\ - \left(\frac{d}{dr} + \frac{\kappa}{r} \right) S_{KK'}(r, \omega_1) - (m + \varepsilon_1 - \omega_1 - V(r)) T_{KK'}(r, \omega_1) &= \frac{3}{\omega_1} L_{KK'}(r, \omega_1) \end{aligned} \quad (22)$$

The inhomogeneous terms $K_{KK'}$ and $L_{KK'}$ and the terms U_K and V_K in the integrand of Eq. (22) are

$$K_{KK'}(r, \omega_1) = \begin{cases} \frac{\kappa - \kappa'}{2} (j_1'(\omega_1 r) + \frac{j_1(\omega_1 r)}{\omega_1 r}) F_2(r) - \frac{j_1(\omega_1 r)}{\omega_1 r} F_2(r) & \text{(Coulomb gauge)} \\ j_1(\omega_1 r) G_2(r) - (\frac{\kappa - \kappa'}{2} + 1) j_2(\omega_1 r) \bar{F}_2(r) & \text{(length gauge)} \end{cases} \quad (23)$$

$$L_{KK'}(r, \omega_1) = \begin{cases} \frac{\kappa - \kappa'}{2} (j_1'(\omega_1 r) + \frac{j_1(\omega_1 r)}{\omega_1 r}) G_2(r) - \frac{j_1(\omega_1 r)}{\omega_1 r} G_2(r) & \text{(Coulomb gauge)} \\ j_1(\omega_1 r) \bar{F}_2(r) - (\frac{\kappa - \kappa'}{2} - 1) j_2(\omega_1 r) G_2(r) & \text{(length gauge)} \end{cases}$$

$$U_K(r, \omega_2) = \begin{cases} - \frac{\kappa + 1}{2} (j_1'(\omega_2 r) + \frac{j_1(\omega_2 r)}{\omega_2 r}) F_1(r) + \frac{j_1(\omega_2 r)}{\omega_2 r} F_1(r) & \text{(Coulomb gauge)} \\ j_1(\omega_2 r) G_1(r) + (\frac{\kappa + 1}{2} + 1) j_2(\omega_2 r) F_1(r) & \text{(length gauge)} \end{cases} \quad (24)$$

$$V_K(r, \omega_2) = \begin{cases} - \frac{\kappa + 1}{2} (j_1'(\omega_2 r) + \frac{j_1(\omega_2 r)}{\omega_2 r}) G_1(r) - \frac{j_1(\omega_2 r)}{\omega_2 r} G_1(r) & \text{(Coulomb gauge)} \\ j_1(\omega_2 r) F_1(r) + (\frac{\kappa + 1}{2} - 1) j_2(\omega_2 r) G_1(r) & \text{(length gauge)} \end{cases}$$

Here, $G_a(r)$ and $F_a(r)$ are the large and small components of the electron wave function, satisfying the radial equation

$$\begin{aligned} (m - \epsilon_a + V(r)) G_a(r) + \left(\frac{d}{dr} - \frac{n}{r} \right) F_a(r) &= 0 \\ - \left(\frac{d}{dr} + \frac{n}{r} \right) G_a(r) - (m + \epsilon_a - V(r)) F_a(r) &= 0 \end{aligned} \tag{25}$$

These formulas are formally identical to those for the hydrogenic case. The only difference is in the potential $V(r)$. In Eqs. (24) and (25), $V(r)$ is the potential due to the nucleus and the other electrons, but we do not exclude the core contribution in the inhomogeneous equations (22). In the hydrogenic case, $V(r)$ is the pure Coulomb potential due to the nucleus. The notation used here is the same as that in Parpia and Johnson's paper.² If we set $K' = -1$ in Eq. (23), we obtain the $s \rightarrow s$ decay rate, which is the same as the formula of Parpia and Johnson.² If we set $K' = 2$, and -3 , we find the $d_{3/2} \rightarrow s$ and $d_{5/2} \rightarrow s$ decay rates.

ACKNOWLEDGMENTS

This work was encouraged by W. R. Johnson. The author thanks B. Crasemann for stimulating discussions and helpful comments on the manuscript. The author acknowledges fruitful discussions with Xiao-Ling Liang, Xing-Dong Mu and T. Aberg. This work was partially supported by the Air Force Office of Scientific Research under Grant 87-0026 and by the National Science Foundation under grant PHY-85 16788.

1. K. Ilakovac, J. Tudoric-Ghemo, B. Busic and V. Horvat, Phys. Rev. Lett. 56, 2469 (1986).
2. W. R. Johnson, Phys. Rev. Lett. 29, 1123 (1972); F. A. Parpia and W. R. Johnson, Phys. Rev. A 26, 1142 (1982).
3. S. P. Goldman and G. W. F. Drake, Phys. Rev. A 24, 183 (1981).
4. V. Florescu, Phys. Rev. A 30, 2441 (1984).
5. T. Aberg, in Atomic Inner Shell Processes, edited by B. Crasemann (Academic, New York, 1975), p.353.
6. I. Freund, Phys. Rev. A 7, 1849 (1973); Y. B. Bannett and I. Freund, Phys. Rev. A 30, 299 (1984).
7. X. Mu and B. Crasemann, Phys. Rev. Lett. 57, 3039 (1986).
8. G. E. Brown, R. E. Peierls, and J. B. Woodward, Proc. R. Soc., Ser. A 227, 51 (1954).
9. V. B. Berestetskii, E. M. Lifshitz and L. P. Pitaevskii, Quantum Electrodynamics, (2nd Ed., p. 482., 1982).
10. E.J. Robinson, Phys. Lett. A 117, 465 (1986).
11. M. Crance and Mireille Aymar J. Phys. B20. L155 (1987).


FIGURES CAPTIONS

Fig. 1. Feynman diagrams for advanced process


(a) in terms of a hole, (b) in terms of an electron.

Fig. 2. Feynman diagrams for retarded process

(a) in terms of a hole, (b) in terms of an electron.




(a)




(b)

Fig. 1



(a)



(b)

Fig. 2

END

FEB.

1988

DTIC